

Ant Killer



Plus Multi-Purpose Insect Control₂

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Keep children and pets off treated area until product has been watered in and lawn is dry

RESTRICTIONS:

- Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of runoff.
- Do not make applications during rain.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
 Do not apply this product to edible growing crops or stored raw agricultural commodi-
- ties used for food or feed.
- Do not allow applications to contact water inhabited by fish, such as in ornamental fish ponds that are located around structures being treated.
- Do not apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, do not apply this pesticide within 25 ft. of lakes, reservoirs, rivers, perma-
- nent streams, marches or natural ponds, estuaries, and commercial fish farm ponds.

 In New York State, this product may not be applied within 100 feet of a coastal marsh, or stream that drains directly into a coastal marsh.
 Observe all restrictions and precautions which appear on the label of this product.
- Do not apply more than 0.36 lbs. of the active ingredient (800 pounds of this product) per acre per year.

IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in outdoor ornamental gardens or parks, or on lawns or grounds.

GENERAL INFORMATION

For use outdoors, on lawns and as a band treatment around house foundation.

Kills ants (including Fire Ants), crickets, earwigs, fleas, roaches (German and American) and other insect pests listed in the tables below. To use this product follow these 2 easy steps – (1) Determine the size of the job and measure the area to be treated. (2) Treat as directed in rate chart. Complete coverage is important. Water in granules immediately following application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. PESTICIDE STORAGE: Store in a cool, dry area which is inaccessible to children and pets. Reclose container tightly after each use. **PESTICIDE DISPOSAL**: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **SPILLS**: If the container is leaking or material is spilled, carefully sweep up spilled product and wash exposed area. In spill or leak incidents, keep unauthorized people away.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides.

First Aid: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Poison Control Center for emergency medical treatment at 1-800-222-1222. IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person. IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS: This product is highly toxic to fish and aquatic organisms.

To protect the environment, do not allow pesticide to enter or run off into storm drains, To protect the environment, do not allow pesticute to enter or full of into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind and rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

Note: Buyer assumes all responsibilities for safety and use not in strict accordance with directions

Treatment around Foundations / Tratamiento alrededor de cimientos

Use Instructions Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance), Fleas, Apply at the rate of 2 lb. to 3 lb. per 1,000 sq. ft. Apply a 3 to 10 foot band of granules next to the house foundation and completely distribute the granules. (an area 5'x 200') or 3.2 oz. to 4.8 oz. per 100 Water lightly immediately following application. Treat when insects first appear or when you notice lawn Roaches (German and American), Palmetto bugs, Waterbugs, Millipedes, Mites, sq. ft. (10' x 10'). damage. For best results, leaf litter and debris should be removed from the foundation area. Branches and Silverfish, Sowbugs (Pillbugs), Crickets, Earwigs vegetation that touch the foundation should be cut back. Other potential pest harborage sites, such as deep mulch, should be raked away before application. **Plagas** Proporción de uso Instrucciones de uso Hormigas (argentinas, sureñas, del campo, de Allegheny, carpinteras de Florida, carpinteras negras, del maíz, de la miel, del pavimento, hormiga roja de fuego Aplique a una proporción de 2 libras a 3 libras por cada 1.000 pies cuadrados (0,91 kg a 1,36 kg Aplique una franja de 3 a 10 pies (0,91 m a 3 m) de gránulos cerca del cimiento de la casa y distribuya completamente los gránulos. Riegue ligeramente de inmediato luego de la aplicación. Trate cuando los insectos aparezcan por primera vez o cuando note que hay daño en el césped. Para obtener los mejores importada, molestas, pulgas, cucarachas (alemanas y americanas), de la palma, por 92,9 m²) (un área de 5 pies x 200 pies) (1,5 m x 61 m) ó 3,2 onzas a 4,8 onzas por cada periplaneta americana, ciempiés, ácaros, lepisma, cochinilla de humedad resultados, la hojarasca y los desechos se deben retirar del área de los cimientos. Se debe cortar la 100 pies cuadrados (3,2 onzas a 4,8 onzas por 9,3 m²) (10 pies x 10 pies) (3 m x 3 m). vegetación y las ramas que rocen los cimientos. Antes de la aplicación, se deben barrer otros posibles

| | 3,3 III) (TO pies x TO pies) (3 III x 3 III). | | lugares de allidación de piagas, como compuestos organicos profundos. |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Spot Treatment / Tratamiento tópico | | | |
| Pests | Use Rate | Use I | nstructions |
| Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance) | Mound Treatment: Apply 1–2 teaspoons over and around each ant hill. | Water lightly immediately following application. NOTE: Use both broadcast and spot treatment for difficult to large ant infestations. If soil is dry, water lightly before application to bring the ants closer to the surface for better control. | |
| Fire Ants | Mound Treatment: Apply 1/2 cup over, and 2 ft. around, each fire ant mound. | To prev | diately following application, water in completely. Apply a minimum of 1 gallon of water to each mound area. went the disturbance of ants, water gently. Use of high pressure watering equipment may disturb the ants and migration, reducing product effectiveness. NOTE: For best results, apply in cool weather, 65°-80°F, or in early ng or late evening hours. Treat new mounds as they appear. Fast acting. Works within 24 hours. |
| Plagas | Proporción de uso | Instru | ucciones de uso |
| Hormigas (argentinas, sureñas, del campo, montoneras de Allegheny, carpinteras de Florida, carpinteras negras, del maíz, de la miel, del pavimento, hormiga roja de fuego importada, molesta) | Tratamiento de hormiguero: Aplique 1 a 2 cucharaditas sobre y alrededor de cada hormiguero. | para pl | s ligeramente inmediatamente después de la aplicación. NOTA: Use el tratamiento por dispersión y el localizado lagas de hormigas difíciles o grandes. Si la tierra está seca, riéguela ligeramente antes de la aplicación para ar a las hormigas a la superficie y así obtener un mejor control. |
| Hormigas del fuego | Tratamiento por hormiguero: Aplique 1/2 taza sobre y 2 pies (60 cm) alrededor de cada hormiguero. | área de | e completamente inmediatamente en seguida de la aplicación. Aplique un mínimo de 1 galón de agua a cada le hormiguero. Riegue suavemente para evitar que las hormigas se alboroten. El uso de equipo de riego con lesión puede alborotar las hormigas y ocasionar que emigren, disminuyendo la efectividad del producto. |

ACTIVE INGREDIENT/INGREDIENTE ACTIVO: Lambda-cvhalothrin 0.04% OTHER INGREDIENTS/OTROS INGREDIENTES: 99.96% TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN

NOTA: Para mejores resultados, aplique agua fría 65°-80° (18°-27° C) o en las horas tempranas de la mañana, o tarde en la noche. Trate los nuevos hormiqueros a medida que aparezcan. Actúa rápido, en el lanso de 24 horas

> MANTENGASE FUERA DEL ALCANCE DE LOS NIÑOS **PRFCALICIÓN**