



#382KETTLECLN

Kettle Kleen Decarbonizer

Manufactured for:
Carnival King Supplies

Lancaster, PA 17602

info@carnivalkingsupplies.com



General Description

KETTLE KLEEN is designed to remove carbon deposits from sheet pans, cookware, oven and stove parts and racks. KETTLE KLEEN dissolves the backed on carbon deposits without scraping or scrubbing. Safe to use on stainless steel, and plastic. Will not harm coating on sheet pans, bread or roll pans etc.

A white, free flowing granular powder made up of alkaline cleaning compounds, wetting agents, inhibitors and suspending agents. Not safe on painted surfaces, magnesium or zinc.

Physical Properties

Chemical Composition

Destaining agents, alkaline builders, surfactants, glaze and metal protectants, anti-caking and water softening agents.

Appearance

White powder

Caustic

Not Present

Metal Safety

Recommended for use on aluminum and other soft metal ware at normal use concentrations.

Foaming

Low

Rinsing

Excellent

Water Hardness Tolerance

10 Grains Per Gallon at 0.3%

7 Grains Per Gallon at 0.2%

3 Grains Per Gallon at 0.1%

Packaging

31 oz container - 2 lbs net

Recommended Range of Use Concentrations

Soak Tank

Use plastic drum or metal tank. Make up solution with warm water using 8 pounds in 25 to 30 gallons of water. Immerse carbonized pots, pans, sheet pans, racks and stove parts in solution at room temperature. Allow to soak, preferably overnight. Thoroughly rinse, with light brushing where needed.

Spray Washer

Use 1 to 4 ounces of KETTLE KLEEN per gallon of hot water. Temperature should be 140 °F to 170°F. Always follow with spray rinse using clean water.

Manual Cleaning

Use 1/4 to 1 ounce of KETTLE KLEEN per gallon of warm water (110°F to 130°F). Apply with sponge, brush, hand spray, etc. Follow with clean water rinse.

CAUTION: WARNING KEEP OUT OF REACH OF CHILDREN.

Alkaline powder. Contact with skin and eyes may cause severe irritation. Harmful if swallowed.

First Aid

IF IN EYES: Flood with water if splashed in eyes. Flush with water for 15 minutes and consult physician immediately. IF INGESTED: Drink large amounts of milk, milk of magnesia or gelatin, or if these are not available, drink large amounts of water. DO NOT induce vomiting. Obtain prompt medical attention.